

List of caddisflies (Insecta: Trichoptera) collected by Bulgarian scientists in Turkey

Krassimir KUMANSKI, Füsün SIPAHILER

Abstract. A faunistic list of caddisflies (Trichoptera), based on the material collected from 52 localities in Turkey by Bulgarian scientists is given in the present study. The list contains 67 species belonging to 10 families. *Micropterna fissa* McLachlan is newly recorded in Turkey. The first two trogloxene caddisflies in Anatolia are reported.

Key words: Trichoptera, Turkey, Faunistics

Introduction

The Trichoptera fauna of Turkey was poorly known until publishing the first list in 1984 (MALICKY & SIPAHILER, 1984). In that paper 204 species are listed. A second detailed list was presented in 1987, increasing the total number of known species to 235 (SIPAHILER & MALICKY, 1987). In 1993 and 1995 additional lists were published, so the number of known species increased to 313 (MALICKY & SIPAHILER, 1993; SIPAHILER, 1996). After that, many new species and new records were published, so the number of the known species of Turkey reached 366. In this paper, a faunistic list of Trichoptera is given, based on the material collected for several years mainly by Bulgarian scientists. Almost all of the material was collected on light. The list of the determined taxa, which contains 2671 specimens, includes 67 species belonging to 23 genera and 10 families, respectively (Table 1). Among them *Micropterna fissa* McLachlan, 1875 is newly recorded in Turkey and *Micropterna hatatitla* Malicky, 1974 and *Stenophylax meridiorientalis* Malicky, 1980 are the first regular trogloxene species collected in caves in Anatolia. The list of the localities is arranged by geographic regions (SIPAHILER & MALICKY, 1987).

List of the localities

List of the localities

Region 1 (Marmara Region)

1. Çatalca, Akalan vill., 6.4.1913, leg. D. Iltchev.
2. Adapazari, Reşadiye vill., near Kuzuluk, 300 m, (on light), 7.6.1996, leg. S. Beshkov.

Region 3 (Northwestern Anatolia)

3. Bolu, Gerede, 17 km west of Çerkes, 4 km before Kurt Çimeni, (on light), 11.7.1995, leg. S. Beshkov & J. Gelbrecht¹.
4. Bolu, Çamlık vill. near town of Yeniçaga, (on light), 1200 m, 25.7.1995, leg. S. Beshkov & J. Gelbrecht.

Region 4 (Central Anatolia)

5. Ankara, Kızılıcahamam, Pazar, 1000 m, 20.7.1991, leg. S. Beshkov.
6. Nigde, Nevşehir, Capadoccia, above Uçhisar, on the road to Ürgüp, 1415 m (on light), 13.5.1999, leg. S. Beshkov & J. Gelbrecht.
7. Capadoccia, Nevşehir, Zelve vill. near Göreme, 1110 m, 22.7.1995 (on light), leg. S. Beshkov & J. Gelbrecht.
8. Aksaray, Hasan Dagh, 20 km S of Aksaray, A. Dikmen vill., near Tashpinar (1100 m), Melendcz Mt., (on light), 9.6.1996 leg. S. Beshkov.
9. Nigde, Ürgüp, Capadoccia, 1100m, 16.6.1996, (on light) leg. S. Beshkov.
10. Çankırı, near Ilgaz, 20.6.1996, (on light), leg. S. Beshkov.
11. Ankara, Kızılıcahamam, Kızılıcaköy, 900m, 8.6.1996, (on light), leg. S. Beshkov.
12. Sivas, near Zara, 1570 m, 24.6.2000, (on light), leg. S. Beshkov.

Region 5 (Southern Anatolia)

13. Antalya, Bayır, 750m, 5 km from Güzelbag, (on light), 26.11.1998, leg. S. Beshkov.
14. Antalya, Alanya, above Pasşalar, 250 m, (on light), 25.11.1998, leg. S. Beshkov.
15. Antalya, SW of Antalia, 13 km on the road to Altinyaka to Denizli, 700 m, (on light), 23.11.1998, leg. S. Beshkov.
16. Antalya, Termesos National Park, 25 km from Antalia to Denizli, 600 m, (on light), November, 1998, leg. S. Beshkov.
17. Beyşehir, Dumanlı vill., Düden Yayla, Almalaike cave, 5.7.1993, leg. P. Beron.
18. Beyşehir, Çamlık vill., Balatini cave, 11.7.1993, leg. P. Beron.
19. Akşehir, Sultan Dag, 1700 m, 25-26.6.1996, leg. B. Guéorguiev.
20. Nigde, Çamardi, Aladaglar, 1300 m, river, 10 km S from Çamardi, 27-28.6.1996, leg. B. Guéorguiev.

¹ J. Gelbrecht, T. Drechsel and B. Schacht - German entomologists, partly accompanying the Bulgarian entomologists

21. Akşehir, Sultan Dagi, 1000 m, oak forest, 25.6.1996, leg. B. Guéorguiev.
22. Mersin, Bolkar Dagi, above Çamliyayla, 1600 m, 11.6.1996 (on light), leg. S. Beshkov, J. Gelbrecht & T. Drechsel².
23. Adana, Osmanye, Yarpuz vill., 980 m, 12.6.1996, (on light), leg. S. Beshkov.

Region 6 (Northeastern Anatolia)

24. Artvin, Karadeniz Mts., above Yusufeli, 1800 m, 15.7.1995, (on light), leg. S. Beshkov & J. Gelbrecht
25. Artvin, Karadeniz Mts., under Kaçkar Peak, near Yaylalar vill. above Yusufeli, 1780 m, 17.7.1995, leg. S. Beshkov & J. Gelbrecht
26. Artvin, Karadeniz Mts., above Yusufeli, Bahral vill., 945 m, on the way to Sartgöl, (on light), 16.7.1995, leg. S. Beshkov & J. Gelbrecht.
27. Artvin, Karadeniz Mts., above Yusufeli, 630 m, on the way to Sartgöl, (on light), 20.7.1995, leg. S. Beshkov & J. Gelbrecht.
28. Amasya, Yenice, Direkli vill., 13 km before Amasya from Akdag, 580 m, (on light), 13.7.1955, leg. S. Beshkov & J. Gelbrecht.
29. Erzurum, Tortum, 1580 m, near dried river, (on light), 3-4.7.1996, leg. B. Guéorguiev.
30. Giresun, Giresun Daglari, Egridin pass, 1450 m (on light), 4-5.7.1996, leg. B. Guéorguiev.
31. Erzurum, 10 km southern above Ispir, 1800-2010 m (on light), 25.6.2000, leg. S. Beshkov.
32. Erzurum, Karadeniz Daglari, ca. 6 km southern above Ispir, 1600 m (on light), 26.6.2000, leg. S. Beshkov & B. Schacht².
33. Karadeniz Daglari, ca. 15 km northern of Ovit Dagi Geçidi, 1700 m, (on light), 27.6.2000, leg. S. Beshkov.
34. Karadeniz Daglari, Ovit Dagi, southern slopes, above Ovit Dagi Geçidi, 2700 m, (on light), 29.6.2000, leg. S. Beshkov & B. Schacht.
35. Karadeniz Daglari, Ovit Dagi, northern slopes, above Ovit Dagi Geçidi, 2100 m, (on light), 30.6.2000, leg. S. Beshkov & B. Schacht.
36. Karadeniz Daglari, ca. 10 km south above Ispir, above Köpencuk vill., 1850 m (on light), 4.7.2000, leg. S. Beshkov & B. Schacht.
37. Karadeniz Daglari, Ispir, Düzköy above Çoruh valley, 2200 m (on light), 1.7.2000, S. Beshkov.
38. Karadeniz Daglari, Prov. Erzurum, Çoruh valley, 20 km E from Ispir, on the road to Yusufeli, 1060 m (on light), 22.4.2001, leg. S. Beshkov & J. Gelbrecht.
39. Karadeniz Daglari, near Köprüköy vill., 2300 m (on light), 27.7.2001, leg. S. Beshkov.
40. Karadeniz Daglari, below Ovit Dagi Geçidi, 2400 m (on light), 27.7.2001, leg. S. Beshkov.
41. Erzurum, Ispir, Mescit Daglari, Çoruh valley, near Yukari Ozbar, 1300 m (on light), 7.7.2001, leg. S. Beshkov.
42. Erzurum, Karadeniz Daglari, Çoruh valley, 25 km from Ispir to

² see Footnote 1 on previous page

Yusufeli, 1040 m (on light), 3.8.2001, leg. S. Beshkov.

43. Erzurum, above Köprüköy, Ispir, 1900 m (on light), 4.8.2001, leg. S. Beshkov.

44. Erzurum, Karadeniz Daglari, near Köprüköy vill., 2030 m (on light) 28.7.2001, leg. S. Beshkov.

Region 7 (East Anatolia)

45. Sivas, near Gürün, 1415 m (on light), 21.7.1995, leg. S. Beshkov & J. Gelbrecht.

46. Erzurum, 20 km south of Gölyurt Geçidi, 2000 m (on light), 5.7.2000, leg. S. Beshkov & B. Schacht.

47. Erzurum, southern side of Gölyurt Geçidi, 1900 m (on light), 3.7.2000, leg. S. Beshkov.

48. Erzurum, 25 km from Gölyurt Geçidi, near Kirik vill., 2100 m (on light), 2.7.2000, leg. S. Beshkov.

49. Erzurum, Ispir, above Köprüköy, 1960 m (on light), 4.8.2001, leg. S. Beshkov.

50. Erzurum, Ispir, above Köprüköy, 1665 m (on light), 1.8.2001, leg. S. Beshkov.

51. Van, ca. 7 km E of Güzeldere Geçidi, on the road to Bagkale, 2450 m (on light), 31.7.2001, leg. S. Beshkov.

52. Erzurum, ca. 10 km S below Gölnet Geçidi, 2000 m (on light), 5.8.2001, leg. S. Beshkov.

List and localities of the determined taxa

Table 1

List of the determined taxa. Localities - bolded numbers; M - males; F - females; FS - Collection Sipahiler)

Family Rhyacophilidae

Rhyacophila fasciata aliena Martynov, 1916

Rhyacophila nubila Zetterstedt, 1840

Rhyacophila subovata Martynov, 1913

Localities and material collected

26: 1 F; **43:** 1 F

3: 3 M, 3 F; **20:** 1 M; **23:** 9 M, 4 F; **30:** 1 F; **48:**

2 M, 2 F; **50:** 3 F; **52:** 9 M, 4 F

33: 1 F

Family Glossosomatidae

Glossosoma capitatum Martynov, 1913

Agapetus delicatulus McLachlan, 1884

Agapetus sp.

30: 1 F; **50:** 1 M, 1 F

3: 1 M; **23:** 1 M

35: 1 F

Family Hydropsychidae

Allotrichia vilnensis Raciecka, 1937

Hydropsyche forcipata Eaton, 1873

Hydropsyche occulta Eaton, 1873

41: 1 F (in coll. FS)

51: 1 M (in coll. FS)

51: 3 M (1 M in coll. FS)

Family Philopotamidae

Philopotamus sp.

38: 1 F; **43:** 1 F

Family Psychomyiidae

Psychomyia pusilla Fabricius, 1871

11: 5 F; **27:** 1 M; **31:** 1 F; **32:** 2 M; **36:** 1 F;

41: 6 M, 4 F; **42:** 3 F; **43:** 2 F; **44:** 1 M; **45:** 9 M,

12 F; 47: 12 M, 1 F; 48: 12 M, 24 F; 50: 2 F; 52:
20 (M + F)

13: 1 F; 41: 1 F; 43: 2 F

Tinodes sp. (Females)

Family Polycentropodidae

Plectrocnemia latissima Martynov, 1913 43: 1 M

Polycentropus flavomaculatus Pictet, 1884 52: 1 M

Family Hydropsychidae

Cheumatopsyche lepida Pictet, 1884 11: 1 M, 1 F; 27: 20 M, 34 F; 41: 6 M, 13 F;
48: 1 M

Cheumatopsyche capitella Martynov, 1927 23: 2 M; 52: 1 M, 1 F

Hydropsyche cetibeli Malicky et Sipahiler, 1993 13: 1 M

Hydropsyche instabilis Curtis, 1834 2: 2 M; 37: 3 M, 6 F; 41: 1 M; 42: 1 M;
50: 10 M (4 M in coll. FS)

Hydropsyche kebab Malicky, 1974 28: 2 M; 30: 1 M; 46: 4 M; 51: 2 M
(1 M in coll. FS); 52: 89 M (10 M in coll. FS)

Hydropsyche krassimiri Malicky, 2001 11: 1 M (in coll. FS)

Hydropsyche lepnevae Botosaneanu, 1967 3: 5 M; 27: 1 M; 31: 1 M; 32: 23 M, 5 F;
34: 42 M, 15 F; 37: 89 M, 14 F; 43: 2 M;

Hydropsyche mahrkusha Schmid, 1959 46: 2 M; 48: 1 M; 50: 8 M (6 M in coll. FS); 52: 2 M

Hydropsyche martynovi Botosaneanu, 1967 25: 1 F; 40: 1 F

Hydropsyche orduensis Sipahiler, 1987 3: 12 M, 42 F; 50: 1 M

Hydropsyche acuta Martynov, 1909 24: 1 M

Hydropsyche bittis Malicky, 1986 30: 108 M; 43: 1 M

Hydropsyche bulbifera McLachlan, 1878 23: 1 M

Hydropsyche consanguinea McLachlan, 1884 5: 8 M, 6 F; 11: 3 M, 2 F
29: 2 M (1 M in coll. FS); 32: 1 M; 42: 58 M;
44: 7 M, 8 F; 49: 64 M, 58 F; 50: 5 M

Hydropsyche contubernalis McLachlan, 1865 2: 1 M; 3: 1 M; 7: 9 M, 8 F; 28: 3 M; 32: 3 M;
41: 22 M, 6 F; 50: 6 M (3 M in coll. FS)

Hydropsyche cornuta Martynov, 1909 27: 25 M; 28: 1 M; 41: 1 M; 47: 1 M

Hydropsyche modesta Navas, 1925 2: 1 M; 28: 2 M

Hydropsyche cf. Incognita Pitsch, 1993 2: 1 M; 3: 1 M, 22 F; 24: 72 M, 61 F;
26: 62 M, 48 F; 27: 12 M, 42 F; 30: 52 M, 47 F;
32: 1 M, 8 F; 47: 3 M, 3 F

Hydropsyche cf. pellucidula Curtis, 1834 47: 62 M, 48 F

Hydropsyche sakarawaka Schmid, 1959 30: 1 M, 1 F; 33: 4 M, 17 F (2 M, 2 F in coll. FS)

Family Brachycentridae

Micrasema bifoliatum Martynov, 1925 30: 3 M

Family Limnephilidae

Drusus bayburtii Cakin, 1983

24: 8 F; 25: 61 M, 57 F; 27: 1 F; 30: 14 M;
33: 4 M, 2 F; 40: 2 M, 1 F; 435 M, 4 F

46: 1 F

52: 1 F

36: 1 F

25: 1 F

34: 3 M, 1 F (2 M, 1 F in coll. FS)

4: 1 F; 27: 1 M

9: 1 M

11: 1 M; 46: 1 F

1: 1 M; 8: 1 M, 3 F; 17: 4 M, 4 F; 21: 1 M;
34: 1 M; 37: 1 F

<i>Stenophylax permistus</i> McLachlan, 1895	19: 1 F
<i>Micropterna caesareica</i> Schmid, 1959	12: 1 M; 16: 1 M; 21: 1 M; 31: 1 F; 33: 1 M; 36: 1 F
<i>Micropterna coiffaiti</i> Decamps, 1962	15: 2 M, 1 F; 16: 2 M
<i>Micropterna fissa</i> McLachlan, 1875	13: 2 M, 3 F (1 M, 2 F in coll. FS); 15: 1 M
<i>Micropterna hatatitla</i> Malicky, 1985	12: 2 M, 1 F (1 M, 1 F in coll. FS); 18: 6 M, 22 F; 31: 1 F; 34: 1 M, 2 F
<i>Micropterna nycterobia</i> McLachlan, 1875	19: 4 M, 4 F; 21: 4 M, 1 F
<i>Micropterna sipahilarae</i>	
Kumanski et Malicky, 1997	30: 1 M (the Holotype)
<i>Micropterna taurica</i> Martynov, 1917	6: 1 M; 8: 1 M; 12: 1 F
<i>Micropterna muehleni</i> McLachlan, 1884	12: 1 M, 1 F
<i>Micropterna</i> sp. 1	14: 1 F
<i>Micropterna</i> sp. 2	10: 1 F
<i>Micropterna</i> sp. 3	35: 1 F
<i>Potamophylax luctuosus armeniacus</i> Mey, 1979	24: 4 M, 1 F; 25: 14 M, 2 F; 26: 4 M; 33: 14 M, 4 F; 34: 3 M; 35: 9 M
Family Sericostomatidae	
<i>Sericostoma grusiense</i> Martynov, 1913	24: 14 M, 22 F; 25: 1 F; 26: 7 M, 12 F; 27: 1 M, 2 F; 35: 1 F; 36: 1 F
<i>Sericostoma flavicorne</i> Scneider, 1845	29: 16 M, 4 F (6 M, 2 F in coll. FS); 30: 18 M, 11 F (10 M, 5 F in coll. FS)
Family Lepidostomatidae	
<i>Lepidostoma hirtum</i> Fabricius, 1775	3: 2 F; 47: 1 M (in coll. FS)
<i>Dinarthrum chalдыrense</i> Martynov, 1909	29: 1 F; 51: 1 M
Family Leptoceridae	
<i>Athripsodes iltsci</i> Malicky, 1987	41: 3 M (1 M in coll. FS)
<i>Athripsodes longispinosus</i> Martynov, 1909	3: 2 M, 2 F; 11: 2 M, 1 F
<i>Athripsodes sewangensis</i> Martynov, 1925	52: 14 M, 36 F (6 M, 6 F in coll. FS)
<i>Setodes dehensurae</i> Cakin et Malicky, 1983	42: 1 M; 43: 5 M (3 M in coll. FS)
<i>Setodes viridis</i> Fourcroy, 1785	41: 3 M, 1 F; 42: 11 M, 3 F (4 M, 1 F in coll. FS)

Remarks

***Hydropsyche cetibeli* Malicky et Sipahiler.** This species was described occurring in SW Turkey, Cetibeli (Province of Mugla) (MALICKY & SIPAHILER, 1993, and found further on in several localities in the Province of Antalya (Murtiçi, Akseki, Manavgat, Cayi) (SIPAHILER, unpublished data). It is an endemic of the SE-Mediterranean region, known outside Turkey only on some Greek E-Aegean islands - Chios, Lesbos and Samos (MALICKY, 2001).

***Hydropsyche ordensis* Sipahiler.** This endemic species of NE Anatolia was described occurring in the Province of Ordu (SIPAHILER, 1987). Later on it was reported from Çavşat, Çermik Köyü (Province of Artvin) (SIPAHILER, 2001). According to latest data (Loc. 24 in the present paper), the species occurs also in the Province of Rize (Kaçkar Mountains).

***Hydropsyche bitlis* Malicky.** This SE Anatolian endemic was recorded and described in the eastern Province of Bitlis (MALICKY, 1986), and later reported from Saimbeyli (S Anatolia, the northern part of the Province of




Fig. 1. *Micropterna* sp. 1, Female genitalia: 1 - lateral view; 2 - dorsal view; 3 - ventral view

Adana) (SIPAHILER, 1996). The Province of Adana is probably the westernmost border of its distribution.

Hydropsyche cf. incognita Pitsch. and *Hydropsyche cf. pellucidula* Curtis. Because of some still existing taxonomic problems within the *pellucidula*-species group (especially in the Anatolian region) these two species cannot be considered definitely determined.

Micropterna hatatitla Malicky. This species was recorded and described in the Province of Ankara (MALICKY, 1985) and considered an endemic of Anatolia. Further on, it was found in several southern (Antalya, Elmali, Gömbe /2800 m, 26.06.1988, 1 M and 1 F pupae/, Konya, Hadim, Geyik Dagi, Egrigöl /2000, 01.09.1996, 8 M, 2 F pupae/) and north-western regions




Fig. 2. *Micropterna* sp. 2, Female genitalia: 1 - lateral view; 2 - dorsal view; 3 - ventral view

of Turkey (Kasatmonu, 14 km south of Kûre, 1200 m, 02.06.2002, 6 M and 2 F pupae) (SIPAHILER, unpublished data). The additional data in the present paper show that it is broadly distributed also in NE Turkey (Locs. 12, 31, 34). Locality Nr. 18 in the Taurus Mts. near Beysehir is the first report of a Trichoptera found in a cave in Turkey. However, the species was discovered also in the E Rhodopes Mts., Bulgaria (KUMANSKI, 1993), as well as in the Pirin Mts. in the same country (Bayovi Dupki Circus, cave No. 29, 22.09.2002, 1 M, leg. P. Beron & B. Petrov) (KUMANSKI, unpublished data). Thus, *M. hatatitla* should be considered a SE European-Anatolian species rather than an endemic of Turkey.




Fig. 3. *Micropterna* sp. 3, Female genitalia: 1 - lateral view; 2 - dorsal view; 3 - ventral view

***Micropterna* spp. 1, 2, 3.** These females belong to three different taxa and have not been described so far. They (or, some of them) could belong to either already known species, modeled after males, or to new ones. Their species association so far seems rather preliminary, therefore we give only their genital figures (Figs. 1, 2 and 3) and no other comments.

Potamophylax luctuosus armeniacus Mey. According to the opinion of one of us (KK) this taxon could be elevated to a specific status. However, a revision of the whole material available (including the type series of *P. l. armeniacus* (MEY, 1979)) needs to be done to solve the problem. We would only note, that all the males mentioned here do not show any trace of variability of the genital structures. A similar revision is to be done soon (MEY & KUMANSKI, unpublished data).

Setodes dehensure Cakin et Malicky. This is an endemic species of Turkey. It was recorded and described in S Anatolia (Antalya, Manavgat, Köprüçay), and later on discovered further eastwards in the Province of Antalya (Gündogmuş, Alara Çayı) and the Province of Adana, Saimbeyli (SIPAHILER, unpublished data). Now it was found also in the NE Turkey (Locs. 42 and 43).

Acknowledgement

We wish to thank the Scientific Research Council of Turkey (TUBITAK-BAYG NATO-D), for supporting this research.

References

- KUMANSKI K. 1993. Addition to Volume 15 (Trichoptera, Annulipalpia) and Volume 19 (Trichoptera, Integripalpia) of the Series "Fauna of Bulgaria". - Hist. nat. bulgarica, 4: 39-64. (In Bulgarian).
- MALICKY H. 2001. Ein Beitrag zur Kenntniss der Arten der *Hydropsyche instabilis*-Verwandtschaft im östlichen Mittelmeergebiet (Trichoptera, Hydropsychidae). - Linzer biol. Beitr., 33 (1): 489-518.
- MALICKY H., SIPAHILER F. 1984. A faunistic survey of the caddisflies (Trichoptera) of Turkey. - In: Morse J. (ed.) Fourth. Int. Sym. on Trichoptera. Ser. Entomologica, Junk, 24: 207-212.
- MALICKY H., SIPAHILER F. 1993. Köcherfliegen (Trichoptera) aus der Türkei, mit Bemerkungen zu weiteren Mittiterranen Köcherfliegen. - Mitt. Schweiz. Ent. Ges., 66: 457-478.
- MEY W. 1979. Die *Potamophylax luctiosus* Piller - Verwandschaftsgruppe in Kaukasus (Trichoptera, Limnephilidae). - D.E.Z., N.F., 26 (1-3): 39-41.
- SIPAHILER F. 1987. Türkiye'deki *Hydropsyche* cinsi *instabilis* grubu (Trichoptera, Hydropsychidae) erkeklerinin sistematik yönünden incelenmesi. - DOGA, TU Zooloji D.C., 11 (3): 161-178.
- SIPAHILER F. 1996. Studies on the Trichoptera fauna of southern Anatolia. - Entomofauna (Anselfelden), 17 (16): 293-309.
- SIPAHILER F. 2001. Camili bölgesinin faunistik özelliklerinin incelenmesi (Trichoptera, Insecta). - Kırsal Çevre Yıllığı, 2000: 81-88.
- SIPAHILER F., MALICKY H. 1987. Die Köcherfliegen der Türkei (Trichoptera). - Entomofauna, 8: 77-165.

Received on 18.11.2002

Authors' addresses:

Dr Krassimir Kumanski
National Museum of Natural History
Tsar Osvoboditel Blvd. 1
1000 Sofia, Bulgaria
E-mail: kumanskik@yahoo.com

Prof. Dr Füsün Sipahiler
Hacettepe Universitesi
Egitim Fakültesi
Biyoloji Egitimi Anabilim Dali
Beytepe, Ankara, Turkey

Списък на ручейниците (Insecta: Trichoptera), събрани от български зоолози в Турция

Красимир КУМАНСКИ, Фюзун СИПАХИЛЕР

(Р е з ю м е)

Съобщавам се фаунистични данни за 67 вида ручейници (Trichoptera), събиранни в Турция през периода 1993-2000 г. предимно от български зоологи. С изключение на специално упоменатите екземпляри, материалът се съхранява в алкохолните колекции на Националния природонаучен музей в София. *Micropterna sipahileri Kumanski et Malicky* е описан като нов за науката в друга публикация. Видът *Micropterna fissa McLachlan* е нов за фауната на Турция. *Stenophylax meridiorientalis McLachlan* и *Micropterna hatatitla Malicky* са първите ручейници, съобщени от турски пещери. Те принадлежат към категорията "редовни троглоксени" ("regular trogloxenes"). *M. hatatitla* е същевременно нов член от тази екологична категория. Три други вида от ред *Micropterna* са установени само по женски индивиди. Тъй като засега не може да бъде уточнено дали те (или някои от тях) принадлежат към нови за науката видове, или към видове, чиито женски са неописани до този момент, тук се ограничаваме само с представянето на съответните генитални фигури (фиг. 1, 2 и 3). За някои недостатъчно познати или все още проблематични таксони са направени съответните бележки.